



Revista Mexicana de Biodiversidad

ISSN: 1870-3453

falvarez@ib.unam.mx

Universidad Nacional Autónoma de México  
México

Domínguez-Domínguez, Omar; Pérez-Rodríguez, Rodolfo; Doadrio, Ignacio  
Morphological and genetic comparative analyses of populations of *Zoogoneticus quitzeoensis*  
(Cyprinodontiformes:Goodeidae) from Central Mexico, with description of a new species  
Revista Mexicana de Biodiversidad, vol. 79, núm. 2, diciembre, 2008, pp. 373-383  
Universidad Nacional Autónoma de México  
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=42511935011>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

[redalyc.org](http://redalyc.org)

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



## Morphological and genetic comparative analyses of populations of *Zoogoneticus quitzeoensis* (Cyprinodontiformes:Goodeidae) from Central Mexico, with description of a new species

### Análisis comparativo morfológico y genético de diferentes poblaciones de *Zoogoneticus quitzeoensis* (Cyprinodontiformes:Goodeidae) del Centro de México, con la descripción de una especie nueva

Domínguez-Domínguez Omar<sup>1\*</sup>, Pérez-Rodríguez Rodolfo<sup>1</sup> and Doadrio Ignacio<sup>2</sup>

<sup>1</sup>Laboratorio de Biología Acuática, Facultad de Biología, Universidad Michoacana de San Nicolás de Hidalgo, Fuente de las Rosas 65, Fraccionamiento Fuentes de Morelia, 58088 Morelia, Michoacán, México

<sup>2</sup>Departamento de Biodiversidad y Biología Evolutiva, Museo Nacional de Ciencias Naturales, José Gutiérrez Abascal 2, 28006 Madrid, España.

\*Correspondent: odoming@jupiter.umich.mx

**Abstract.** A genetic and morphometric study of populations of *Zoogoneticus quitzeoensis* (Bean, 1898) from the Lerma and Ameca basins and Cuitzeo, Zacapu and Chapala Lakes in Central Mexico was conducted. For the genetic analysis, 7 populations were sampled and 2 monophyletic groups were identified with a genetic difference of  $D_{HKY} = 3.4\%$  (3-3.8%), one being the populations from the lower Lerma basin, Ameca and Chapala Lake, and the other populations from Zacapu and Cuitzeo Lakes. For the morphometric analysis, 4 populations were sampled and 2 morphotypes identified, 1 from La Luz Spring in the lower Lerma basin and the other from Zacapu and Cuitzeo Lakes drainages. Using these 2 sources of evidence, the population from La Luz is regarded as a new species *Zoogoneticus purhepechus* sp. nov. The new species differs from its sister species *Zoogoneticus quitzeoensis* in having a shorter preorbital distance ( $PrOL/SL \bar{x} = 0.056$ ,  $SD = 0.01$ ), longer dorsal fin base length ( $DFL/SL \bar{x} = 0.18$ ,  $SD = 0.03$ ) and 13-14 rays in the dorsal fin. The new species differs from both members of its sister taxon (*Zoogoneticus tequila* and *Z. quitzeoensis*) at 10 fixed nucleotide positions in the cytochrome *b* gene. We have determined that *Zoogoneticus purhepechus* is distributed in the lower Lerma, upper Ameca, Armeria and Santiago river basins, and Chapala Lake. This new species should be considered endangered of extinction according to the criteria of the MER (Aii,Bi,Ci,Di) and for the IUCN (A-1,b,c,e).

Key words: *Zoogoneticus*, cytochrome b, new species, Mesa Central, Mexico, morphometry.

**Resumen.** Se realizó un estudio genético y morfométrico en poblaciones de *Zoogoneticus quitzeoensis* (Bean, 1898) pertenecientes a las cuencas de los ríos Lerma y Ameca y los lagos de Cuitzeo, Zacapu y Chapala en el centro de México. Para el análisis genético se analizaron 7 poblaciones, identificándose 2 grupos monofiléticos bien diferenciados, con distancias genéticas entre ellos de  $D_{HKY} = 3.4\%$  (3-3.8%), uno de los grupos se distribuye por las cuencas de los ríos Ameca y bajo Lerma y en el lago de Chapala, mientras que el otro incluye las poblaciones de los lagos de Zacapu y Cuitzeo. Se emplearon 4 poblaciones para los análisis morfométricos identificándose 2 morfotipos, 1 de la localidad del manantial La Luz en la cuenca del bajo Lerma y el otro a los lagos de Zacapu y Cuitzeo. Con estas 2 fuentes de evidencia, la población de La Luz es considerada como una nueva especie *Zoogoneticus purhepechus* n. sp. La especie nueva difiere de su especie hermana, *Z. quitzeoensis* por tener una distancia preorbital más corta ( $PrOL/SL \bar{x} = 0.05 - 0.06$ ), la base de la aleta dorsal más larga ( $DFL/SL \bar{x} = 0.17 - 0.20$ ) y presentar entre 13 y 14 radios en la aleta dorsal. La especie nueva difiere de las 2 especies descritas en el género (*Zoogoneticus tequila* y *Z. quitzeoensis*) en 10 posiciones nucleotídicas fijadas para el gen citocromo *b*. *Zoogoneticus purhepechus* se distribuye por las cuencas de los ríos Ameca, Armería, Santiago y bajo Lerma, así como en el lago de Chapala. *Z. purhepechus* debe ser considerada en peligro de extinción de acuerdo a los criterios del MER (Aii,Bi,Ci,Di) y de la UICN (A-1,b,c,e).

Palabras clave: *Zoogoneticus*, cytochrome b, Mesa Central, México, morfometría.

#### Introduction

The Mesa Central of Mexico is characterized by its high diversity of freshwater fishes (Barbour, 1973; Echelle

and Echelle, 1984; Domínguez-Domínguez et al., 2005). A total of 100 native species have been reported, of which 70% are endemic (Guzmán-Arroyo, 1994). This important biological diversity has been attributed to the complex geological and zoogeographic history of central Mexico (Miller and Smith, 1986; Domínguez-Domínguez et al., 2006a). Of the endemic fish fauna of the Mesa Central, the cyprinodontiform fish subfamily Goodeinae (family Goodeidae) is one of the most diverse and interesting.

The subfamily exhibits internal fertilization, matrotrophy and viviparity (Parenti, 1981; Grudzien et al., 1992). When the genus *Zoogoneticus* Meek, 1902, was described, the 14 species of goodeines were included in the Poeciliidae, which also comprised the presently recognized families Profundulidae, Fundulidae, Rivulidae, Cyprinodontidae, and Anablepidae (sensu Parenti, 1981). Meek (1904) placed *Fundulus robustus* Bean 1892, *Platypoecilus quitzeensis* Bean, 1898 and *Fundulus dugesii* Bean, 1887 in *Zoogoneticus*, and simultaneously described 2 new species (*Z. diazi* Bean, 1887 and *Z. miniatus* Bean, 1887).

Regan (1908) proposed the synonymy of *Z. miniatus* with *Z. diazi* and *Z. maculatus* Regan 1904 with *Z. robustus* (Bean, 1892). The revision by Hubbs and Turner (1939), based on the anatomy of the ovary and the trophotaeniae, restricted the genus to include only *Z. quitzeensis* (Bean, 1898), removing other taxa to what are presently 3 different genera of goodeines (*Allotoca* Hubbs and Turner, 1937, *Allophorus* Hubbs and Turner, 1937 and *Allodontichthys* Hubbs and Turner, 1937). Based on molecular characters the genus *Zoogoneticus* is currently placed in the tribe Chapalichthyini (sensu Doadrio and Domínguez, 2004). The genus is currently comprised of the species *Z. quitzeensis* and *Z. tequila* Webb and Miller, 1988. *Zoogoneticus quitzeensis* is widely distributed in Central Mexico whereas *Z. tequila* has a restricted distribution. The former is considered endangered and the second has been considered extinct in the wild (Espinosa-Pérez et al., 1993; Webb and Miller, 1998; SEMARNAT 2002), although a small and restricted population was recently reported (De la Vega-Salazar et al., 2003). Genetic studies have shown that populations of *Z. quitzeensis* have a geographical structure with a consistently high degree of genetic divergence among populations (Doadrio and Domínguez, 2004; Domínguez-Domínguez et al., 2007).

The causes of the ancient population structure may be explained by several volcanic and tectonic events during the Plio-Pleistocene; the population has been subject to different events of dispersion and vicariance that differ in spatial and temporal scale (Domínguez-Domínguez et al., 2006a). Thus, genetic and morphological differences within and between populations have been observed in other non-goodeid fishes from Central Mexico including

*Poeciliopsis infans* (Woolman, 1894) (Mateos et al., 2002) and the genus *Notropis* Rafinesque 1818 (Schönhuth and Doadrio, 2003).

According to a phylogenetic hypothesis proposed by Doadrio and Domínguez (2004), the westernmost populations of *Z. quitzeensis* are genetically different from populations from the Lake Zacapu and La Mintzita, the latter in the Lake Cuitzeo systems. These results suggest that a morphological and more extensive molecular revision of *Zoogoneticus* needs to be conducted to better establish the taxonomic identity of the different populations of the genus that still exist in nature. These patterns of variation between population in the lower and middle Lerma drainages, with evidence of separated closely related clades, are paralleled by the sister species *Skiffia lermiae* Meek 1902, and *S. multipunctata* (Pellegrin, 1901). It is hypothesized that both groups of species owe their origins to the same vicariant event between 1 and 3.5 Mya (Domínguez-Domínguez et al., 2006a).

The purpose of this study, therefore, is to analyze the morphologic and genetic differences among populations of the genus *Zoogoneticus* and to provide the description of a newly recognized species.


## Materials and methods

The study was based on specimens collected using hand and seine nets and electrofishing. All of the sampled specimens were preserved in 70% ethanol. Voucher specimens are housed at the Universidad Michoacana de San Nicolás de Hidalgo (CPUM), the Museo Nacional de Ciencias Naturales de Madrid (MNCN) and the Instituto de Biología, Universidad Nacional Autónoma de México (CNP). The tissues used in genetic analysis are housed at the Museo Nacional de Ciencias Naturales de Madrid (voucher numbers MEX 38, 4271, 4272, 506 and 508).

**Morphological analysis.** Specimens from the type locality (San Cristobal, Cuitzeo Lake, Michoacan) and 3 other populations of *Z. quitzeensis*, (La Luz-Spring, Zamora, Michoacan; La Mintzita Spring, Morelia, Michoacan and Zacapu Lake, Zacapu, Michoacan) were analyzed (Table 1 and Fig. 1). Twenty morphometric characters were measured with digital calipers (0.01 mm) and 4 meristic variables were recorded using a stereoscopic microscope. The abbreviations used for morphometric variables are: SL, standard length; HL, head length; PrOL, preorbital length; ED, eye diameter; InOW, interorbital width; BD, body depth; BLD, body least depth; PAD, pelvic-anal fin distance; PDD, pelvic-dorsal fin distance; PODE, pelvic fin origin to dorsal fin posterior extent distance; DAD, dorsal-anal fin distance; DOAE, dorsal fin origin to anal fin

**Table 1.** Localities and sample sizes for *Zoogoneticus* spp. populations used for morphometric, meristic and genetic analysis. \*Correspond to the type locality for *Z. quitzeoensis*. \*\* Correspond to the type locality for *Z. tequila*. \*\*\*Correspond to the type locality for *Z. purhepechus* sp. nov. +Correspond to sequences obtained from Genbank

Locality	<i>Zoogoneticus</i> spp Drainage	Morphometrics		Meristics		Genetics
		Males	Females	Males	Females	
1. *San Cristobal Spring	Cuitzeo Lake	5	11	5	11	2
2. La Mintzita Spring	Cuitzeo Lake	16	16	16	16	1+
3. Zacapu Lake	Middle Lerma River	11	7	11	7	2+
4. ***La Luz Spring	Lower Lerma River	15	14	15	14	1
5. Orandino Spring	Lower Lerma River	----	----	----	----	2
6. Jaripo Stream	Chapala Lake	----	----	----	----	1+
7. ** El Rincon Spring	Ameca River	----	----	----	----	2+



**Figure 1.** Distribution range of *Zoogoneticus quitzeoensis*; open circles correspond to historical occurrence points where specimens have not been collected in the last 5 years, solid circles correspond to sites where specimens were collected in the last 5 years. For *Z. purhepechus*, open boxes correspond to historical occurrence points where specimens have not been collected in the last 5 years, solid boxes correspond to sites where specimens were collected in the last 5 years. Numbers correspond to localities shown in Table 1.

posterior extent distance; DFL, dorsal fin length; DEAO, dorsal fin posterior extent to anal fin origin distance; AFL, anal fin length; AEDE, anal fin posterior extent to dorsal fin posterior extent distance; EDUP, end of dorsal fin-upper extreme of caudal peduncle distance; EDLP, end of dorsal fin-lower extreme of caudal peduncle distance; EAUP, end of the anal fin-upper extreme of caudal peduncle distance; EALP, end of the anal fin-lower extreme of caudal peduncle distance. The abbreviations for meristic characters are: D, dorsal fin rays; A, anal fin rays; P, pectoral fin rays; GR, gill rakers. All the measurements are in millimetres.

A two-way analysis of variance (ANOVA), comparing both morphometric and meristic characters, was conducted to test sexual dimorphism and variation between populations. Burnaby's method was used to correct size effect (Burnaby, 1966; Rohlf and Bookstein, 1987; Doadrio et al., 2002). All analyses were conducted with the corrected matrix. To identify the variables that contributed most to the variability between populations, a principal components analysis (PCA) was conducted using the covariance matrix for morphometric characters and the correlation matrix for meristic characters. The classificatory hypothesis obtained by PCA was tested by a discriminate function analysis (DFA). All analyses were conducted with the statistics packages NTSYS v.2.1 (Rohlf, 2000) and SPSS v.13.0. Both sets of characters, morphometric and meristic, were analyzed independently. Because of sexual dimorphism involved in the morphometric measurements, these were analysed separately for males and females.

**Genetic analysis.** Six sequences of the gene Cytochrome *b* of *Zoogoneticus* spp. from different localities (Table 1 and Fig. 1) were obtained from GenBank (AF510751-AF510755 and AF510757) and 1 was obtained for the outgroup (*Xenophorus captivus* (Hubbs, 1924), AF510758). The other sequences (including specimens from the type locality of the new species and specimens from the type locality for *Z. quitzeensis*) were obtained using the following protocol. Total cellular DNA was isolated from tissues by a standard proteinase K and phenol/chloroform extraction method (Sambrook et al., 1989). Two overlapping fragments of the cytochrome *b* gene (total of 1140 bp) were amplified via polymerase chain reaction (PCR) for each individual DNA sample.

The primers used for cytochrome *b* in all species were those discussed in Machordom and Doadrio (2001). The amplification process was conducted as follows: 94 °C (2 min), 35 cycles at 94 °C (45 s), 48 °C (1 min), 72 °C (90 s), and 72 °C (5 min). PCR mixtures were prepared in 25 µl reactions with a final concentration of 0.4 µM of each primer, 0.2mM of each dNTP, 1.5mM MgCl<sub>2</sub>, and 1U of Taq DNA polymerase (Biotools). PCR products were checked on 1.5% agarose gels, and cloned using the

pGEM-T vector (Promega) into *Escherichia coli* JM109. Positive clones were sequenced using the Big Dye Deoxy Terminator cycle-sequencing kit (Applied Biosystems). DNA sequences of both strands were obtained using M13 universal (forward and reverse) sequencing primers. All samples were sequenced on an Applied Biosystems 3700 DNA sequencer following manufacturer's instructions. Chromatograms and alignments were visually checked.

The model of DNA substitution that best fitted the data set was selected using Modeltest 3.7 (Posada and Crandall 1998) using the Bayesian information criterion (BIC). The aligned data were analysed with the Bayesian inference method with the program Mr. Bayes 3.1.1 (Hueselsenbeck and Ronquist, 2001) by simulating a Markov chain for 1,000,000 cycles. Based on the HKY+G model obtained by Modeltest, a genetic distance between the 2 groups was obtained using the program Sequencer 6.1.0 (written by B. Kessing and available at <http://nmg.si.edu/>).

## Results

**Morphometrics.** Analysis of variance showed significant differences ( $\leq 0.05$ ) between species for most of the morphometric variables mainly due to differences in standard length, except for the dorsal-fin origin to end of the anal fin. We inferred from this result that the longer dorsal-fin base in *Z. purhepechus* n. sp. influences the DOAE measurement 0.32-0.25 ( $\bar{x} = 0.29$ ) (Table 2). ANOVA for sexual dimorphism showed significant differences in all variables except in the end of the anal-fin to end of the dorsal-fin distance, thus showing that females have a narrower caudal peduncle than males.

The interaction between populations and sexes in most of the morphometric characters do not show significant differences, except for preorbital length. On the contrary, in the meristic characters, the significant differences are only between populations (Table 2). These results justify the separation of morphometric but not meristic analyses by sex.

In an exploratory PCA with the morphometric measurements, PCI explains 90.29% of the variation in males and 90.03% in females; for both sexes, the eigenvectors show closed values with the same symbol, suggesting an influence of the standard length in the results (Doadrio et al., 2002). A second PCA with a Burnaby corrected matrix, accumulates 45.53% of the variance in the PCII in males and 52.42% in females. For males, the high values in eigenvectors were preorbital length, dorsal fin length and anal fin length in the PCI, and interorbital width and FAD-ESPC in the PCII. For females, the high values in eigenvectors were preorbital length, dorsal fin



**Table 2.** Two-way analysis of variance testing for sexual dimorphism, population variation and their interaction (Pop\*Sex). n.s. not significant differences. Mean squares from SigmaStat v. 3.0.1.

Variable	Population	Sex	Pop*Sex
<i>Meristics</i>			
D	0.0210	0.000112n.s.	0.000617n.s.
A	0.000719n.s.	0.000141n.s.	0.000800
P	0.00282	0.0103	0.00259
GR	0.0131	0.000697 n.s.	0.000822 n.s.
<i>Morphometrics</i>			
SL	0.0948	0.139	0.00403n.s.
HL	0.0935	0.0734	0.00715n.s.
PrOL	0.166	0.0761	0.0279
ED	0.0926	0.0408	0.00884n.s.
InOW	0.112	0.0791	0.00903n.s.
BD	0.128	0.103	0.0102 n.s.
BLD	0.0557	0.0376	0.00917 n.s.
PAD	0.0951	0.181	0.0147n.s.
PDD	0.114	0.113	0.00770n.s.
PODE	0.103	0.107	0.00845n.s.
DAD	0.114	0.0547	0.0128n.s.
DOAE	0.121n.s.	0.102	0.0122n.s.
DFL	0.133	0.0502	0.0110 n.s.
DEAO	0.111	0.0855	0.00851n.s.
AFL	0.124	0.000000381n.s.	0.00390 n.s.
AEDE	0.0925	0.139	0.00629n.s.
EDUP	0.0706	0.139	0.00833n.s.
EDLP	0.0839	0.145	0.00514n.s.
EAUP	0.0933	0.148	0.00327n.s.
EALP	0.0838	0.145	0.00314 n.s.

length and FAD-ESPC in the PCI, and anal fin length in the PCII (Table 3). In both sexes, the variation patterns were determined by the preorbital length and dorsal fin length in the PCI. This analysis shows a clear tendency to form 2 groups (ellipses in Fig. 2 A and B), from the lower Lerma basin (La Luz Spring) and middle Lerma (San Cristobal and La Mintzita Springs in the Cuitzeo drainage and Zacapu Lake). With respect to the PCA with meristic characters, PCII accumulates 76.05% of the explained variance, and the high values in the eigenvectors were dorsal fin rays and gill rakers in the PCI, and pectoral fin rays in the PCII (Table 4). Similar to morphometric characters, the meristic characters show a variation pattern with the formation of 2 groups (ellipses in Fig. 3).

Starting with the classificatory hypothesis (formation of 2 groups), the DFA shows a significant difference ( $\alpha \leq 0.05$ ) with the intermediate distances of morphometric characters in males ( $P= 0.003$ ) and females ( $P= 0.010$ ), and with the meristic characters ( $P= 0.000$ ). These results

agree with the proposed classificatory hypothesis and corroborate the variation pattern found in the PCA.

**Genetics.** In the data set, 106 characters were variable, and 35 were parsimony informative. Third codon positions were the most informative characters (24 informative characters), followed by the first codon position (10 characters). Saturation of transition and transversion changes was checked by plotting the absolute number of changes of each codon position against patristic distances. There was no ingroup evidence of saturation at any of the 3 positions (not shown). The HKY-G model was selected as the best fit to the data set. Rate matrix parameters were:  $-\ln L = 2205.0105$ ;  $K = 5$ ;  $BIC = 4456.7324$ . The base frequencies were:  $\text{freqA} = 0.2522$ ;  $\text{freqC} = 0.2685$ ;  $\text{freqG} = 0.1390$ ;  $\text{freqT} = 0.3402$ . Among-site rate variation was approximated with gamma distribution shape parameter  $\alpha = 0.1976$ . The phylogenetic tree obtained by the Bayesian analysis after discarding the initial 500 burn-in chains (Fig. 4) showed the formation of 2 well differentiated groups, with a posterior probability of the branches of 100 for the *Z. quitzeoensis* clade (including the type locality) and 96 for the clade which contains the new species *Z. purhepechus* sp nov. The genetic distance obtained by the model HKY-G within groups was 3.4% (3-3.8%).

### Description

*Zoogoneticus purhepechus* n. sp. (Figures 5 A-B, Table 5)

D = (13) 14; A = 13-14; P = (11-12) 13-15; GR = (7) 9-12. Morphometric measurements are shown in Table 5. Body relatively deep, laterally compressed and elongated, maximum height  $\bar{x} = 3.1$  (range = 2.8-3.6) times the standard length in males and  $\bar{x} = 3.3$  (range = 2.9-3.7) times in females. Minimum body height  $\bar{x} = 6.5$  (range = 6.2-7) times the standard length for males and  $\bar{x} = 7$  (range = 6.2-7.7) in females. Head short, cephalic length  $\bar{x} = 3.5$  (range = 3.3-3.7) times standard length in males and  $\bar{x} = 3.7$  (range = 3.3-4.1) in females. Preorbital distance short, preorbital distance  $\bar{x} = 16.6$  (range = 14.6-21.8) times standard length in males and  $\bar{x} = 20$  (range = 12.8-27.4) in females. Anal fin inserted before origin dorsal fin at same axis. Dorsal fin length long  $\bar{x} = 5.2$  (range = 3.4-7) standard length in males and  $\bar{x} = 5.8$  (range = 5.3-6.6) in females.


**Pigmentation pattern.** When alive, both sexes exhibit a light brown coloration, with dark brown and moderately large spots on the posterior part of the body, starting at the base of the caudal fin. In the anterior part of the body, a

**Table 3.** Eigenvectors and eigenvalues for the first three principal components for 20 morphometric variables in males and females. Variable codes are given in the method section

Eigenvectors	Males			Females		
	I	II	III	I	II	III
SL	0.000	0.004	0.006	0.009	0.001	0.003
HL	0.006	-0.005	0.001	-0.006	0.000	-0.005
PrOL	0.031	-0.028	0.015	-0.073	0.015	-0.003
ED	0.024	-0.016	0.004	-0.013	0.003	-0.017
InOW	0.007	-0.036	-0.007	0.000	0.004	-0.012
BD	0.002	0.004	-0.009	-0.002	-0.008	-0.001
BLD	0.001	0.018	-0.021	0.002	-0.007	0.003
PAD	0.002	-0.009	-0.024	0.017	-0.004	-0.031
PDD	-0.001	0.003	-0.009	0.001	-0.016	-0.006
PODE	-0.007	0.001	-0.004	0.010	-0.016	-0.006
DAD	-0.006	0.001	-0.004	0.006	-0.012	0.006
DOAE	-0.010	0.001	-0.005	0.008	-0.003	0.001
DFL	-0.046	-0.015	0.012	0.025	0.008	0.014
DEAO	-0.011	0.006	0.001	0.005	-0.001	0.000
AFL	-0.024	-0.003	0.015	0.021	0.044	0.004
AEDE	0.008	0.014	-0.009	-0.001	-0.006	0.002
EDUP	0.014	0.022	0.010	-0.026	-0.008	0.015
EDLP	0.009	0.016	0.006	-0.002	-0.007	0.013
EAUP	0.005	0.014	0.016	0.005	-0.004	0.009
EALP	0.011	0.016	0.015	-0.001	0.000	0.017
Eigenvalues	0.005	0.004	0.002	0.008	0.003	0.002
Percentage	24.327	21.206	12.578	37.988	14.435	12.337
Accumulated%	24.327	45.532	58.111	37.988	52.423	64.761

mottling pattern of small spots can be distinguished at the top of the ventral region. They show a pair of dark brown spots laterally aligned at the base of the caudal peduncle, in the region of the hypural plate. In males, during the breeding season, these spots could not be distinguished. The ventral region lacks spots. Adult males are slightly darker than females and may show a slightly bluish or greenish hue on the lateral side of the body and some scales can produce iridescence. The males from the type locality show an intense red band at the end of the pelvic and dorsal fins. In specimens from other localities, this band may be an intense orange. Alcohol preserved specimens are light brown on the body with the abdominal region yellowish, and from the caudal peduncle approximately to the insertion of the anal fin showing moderately large, dark brown spots. From the anal fin forward, spots are smaller and irregularly shaped. The red band in the dorsal and anal fins and the slightly bluish or greenish hue are faint or are less intense.

*Sexual dimorphism and reproduction.* As in all members of the subfamily Goodeinae, males have the first 2 to 7 anal rays shorter. These rays form a short lobe that is inferred to function in sperm transfer (Parenti, 1981) (Fig. 5A). Differences in size were found between sexes, with females being larger than males. As in *Z. quitzeoensis*, the caudal peduncle is narrower (SL/BLD  $\bar{x}$  = 7.0, range = 6.2-7.7) and longer (SL/EDUP  $\bar{x}$  = 4.1, range = 3.4-4.7) in females than in males (SL/BLD  $\bar{x}$  = 6.5, range = 6.1-7.0 and SL/EDUP  $\bar{x}$  = 3.8, range = 3.5-4.4). The males have a stripe with intense red coloration at the end of the dorsal and anal fin. The sex of males can be distinguished at a few weeks after birth. In captive conditions (temperature 21°C  $\pm$  2°C), the males and females can become sexually mature between 8 and 12 weeks. Gestation requires 7 to 9 weeks. Reproduction occurs throughout the year, but peaks when the temperature is between 20 and 21°C. The number of offspring per reproductive event in captivity oscillates between 15 and 45. The fry are usually 9 to 12 mm in



**Figure 2.** Plots of first 2 principal components for 20 morphometric variables corrected by Burnaby's methods for males (A) and for females (B). L) La Luz Spring, C) Cuitzeo Lake (San Cristobal Spring), M) La Mintzita Spring, and Z) Zacapu Lake.

standard length at birth and feed on the second day after birth. First parturition females usually have a low number of offspring.

**Taxonomic summary**

*Material examined. Holotype:* (Table 5, Fig. 5A) CPUM 1509, adult male 34.12 mm SL, La Luz Spring (lower Lerma), Zamora, Michoacán, México, Ludo Couvreur, Jan de Moree, Kees de Jong, Juan C. Merino and Luis Escalera-Vázquez, November 2002.

*Paratypes:* CPUM 1055, (10 individuals); MNCN 246184 (15 individuals); CNP-IBUNAM 14425-14427 (3 individuals). Collected with the holotype.

**Table 4.** Eigenvectors and eigenvalues for the first three principal components for 4 meristic variables. Variable codes are given in the method section

Eigenvectors	I	II	III
D	0.653	0.458	0.405
A	-0.093	0.345	0.176
P	-0.475	0.647	-0.349
GR	-0.905	-0.045	0.457
Eigenvalues	1.479	0.749	0.526
Percentage	50.474	25.578	17.954
Accumulated%	50.474	76.052	94.006


*Distribution:* in accordance with the molecular and morphometric diagnostic characters, we hypothesized that the populations from Ameca, Santiago and Armería basins and Chapala Lake, identified as *Z. quitzeoensis*, should henceforth be considered as *Z. purhepechus*. Thus, the distribution of *Z. purhepechus* occupies the lower Lerma basin, the upper part of the Armería, Santiago and Ameca basins and Chapala Lake (Fig. 1).

*Habitat:* the type locality of *Z. purhepechus* is La Luz Spring. La Luz is a permanent spring of lentic and clear waters and forms a small pond of approximately 1500 m<sup>2</sup>. Once, the water of this spring flowed to the Duero River, which forms part of the lower Lerma River basin. Currently, the water of this spring is used for irrigation and as a water supply to the population. It has an average depth of 1.5 m with a maximum depth of 3.5 m. The bottom is rocky in its periphery and muddy in most of the pond. Aquatic vegetation is *Iris* sp. and *Ceratophyllum* sp, emergent vegetation *Typha* sp. and *Scirpus* sp., and the introduced species *Eichhornia* sp, and terrestrial vegetation is of the subtropical forest type.

The associated fish fauna are the native species, *Allophorus robustus* (Bean, 1892); *Chapalichthys encaustus* (Jordan and Snyder, 1899); *Goodea atripinnis* Jordan 1880; *Skiffia multipunctata* (Goodeidae); *Poeciliopsis infans* (Poeciliidae); and *Lampetra geminis* (Alvarez, 1964) (Petromyzontidae); and the introduced species *Poecilia mexicana* Steindachner 1863; *Xiphophorus hellerii* Heckel 1848, (Poeciliidae); *Oreochromis* spp (Cichlidae); and *Cyprinus carpio* Linnaeus 1758, *Ctenopharingodon idella* (Valenciennes 1844) (Cyprinidae).

*Conservation:* although this species has been taken from a number of localities, and is widely distributed in different drainages along the occidental part of Central Mexico, in the last 5 years, a reduction in its distribution of almost 75% of the historical occurrence points has been






**Figure 3.** Plots of the first 2 principal components for 4 meristic variables. L) La Luz Spring, C) Cuitzeo Lake (San Cristobal Spring), M) La Mintzita Spring, and Z) Zacapu Lake.

observed (Fig. 1). The most common alterations reported in the localities where the species has disappeared are the introduction of exotic species, water pollution and desiccation (De la Vega-Salazar et al., 2003; Domínguez-Domínguez et al., 2005; Domínguez-Domínguez et al., 2006b). Genetic erosion related with human perturbations of the population of this new species was demonstrated recently (Domínguez-Domínguez et al., 2007). This species should be considered as in danger of extinction, following the criteria and categories of the MER-Aii,Bi,Ci,Di (SEMARNAT 2002) and the International Union for the Conservation of Nature and Natural Resources IUCN-A,1a,c,e. (IUCN, 2001-<http://app.iucn.org/webfiles/doc/SSC/RedList/RedListGuidelines.pdf>).

**Etymology:** the name “*purhepechus*” comes from purhepecha, the name of the indigenous ethnic group which inhabited part of the distribution range of this species, including the type locality.

### Remarks

*Zoogoneticus purhepechus* n. sp. differs from its sister species, *Z. quitzeoensis* by the following combination of characters: 13-14 branched rays in the dorsal fin (vs. 12, rarely 11 or 13 branched dorsal rays in *Z. quitzeoensis*), long dorsal fin base length (DFL/SL = 0.18, SD = 0.03 vs.  $\bar{x}$  = 0.16, SD = 0.01 in *Z. quitzeoensis*) and short pre-orbital length (Prol/SL  $\bar{x}$  = 0.056, SD = 0.01 vs  $\bar{x}$  = 0.066, SD = 0.008 in *Z. quitzeoensis*); 2 conspicuous dark brown spots in the hypural plate region, except in males within the reproductive season; 10 molecular autapomorphies




**Figure 4.** Phylogenetic tree of 11 analysed specimens of the genus *Zoogoneticus* recovered from cytochrome *b* sequences (1140 bp) according to the Bayesian analysis based on the best model of evolution that fit our data using the program Modeltest 3.7 (Posada andCrandall 1998). The numbers on the branches represent the Bayesian posterior probability.

in the cytochrome *b* gene also differentiate *Zoogoneticus purhepechus* sp. nov. from *Z. tequila* and *Z. quitzeoensis* (Table 6). Divergences in the cytochrome *b* gene is  $D_{HKY}$  = 3.4% (3-3.8%) compared to *Z. quitzeoensis* and  $D_{HKY}$  = 11% (9-13%) compared to *Z. tequila*.

The genus *Zoogoneticus* is characterized by the presence of some distinctive characteristics, exhibiting 2 to 6 melanic patches in the post-ventral region of the body (Webb and Miller, 1998). Two dark brown spots in the hypural plate region are characteristic, more evident in *Z. quitzeoensis* and *Z. purhepechus* than in *Z. tequila*, in which the spots are less evident or are fused. The trophotaenia is a ribbon-type with 9-14 termini. The genetic and morphometric

**Table 5.** Statistical parameters for morphometric characters in *Z. purhepechus* sp. nov. Each variable is divided by standard length. Variable codes are given in the method section (SD = standard deviation)

Variables	Holotype	Range	14 Males			14 Females		
			Mean	SD	Range	Mean	SD	
SL	34.12	36.95-19.31	28.41	5.45	52.40 - 24.30	33.26	7.08	
HL	0.28	0.30 - 0.27	0.29	0.01	0.30 - 0.24	0.27	0.02	
PrOL	0.08	0.07 - 0.04	0.06	0.01	0.08 - 0.04	0.05	0.01	
ED	0.09	0.10 - 0.08	0.09	0.01	0.10 - 0.07	0.08	0.01	
InOW	0.11	0.12 - 0.09	0.10	0.01	0.11 - 0.10	0.09	0.01	
BD	0.34	0.35 - 0.28	0.32	0.02	0.35 - 0.27	0.30	0.02	
BLD	0.15	0.16 - 0.14	0.15	0.01	0.16 - 0.13	0.14	0.01	
PAD	0.18	0.19 - 0.13	0.17	0.02	0.22 - 0.15	0.18	0.02	
PDD	0.35	0.37 - 0.29	0.34	0.02	0.38 - 0.27	0.32	0.03	
PODE	0.40	0.41 - 0.34	0.38	0.02	0.40 - 0.34	0.37	0.02	
DAD	0.32	0.35 - 0.27	0.32	0.02	0.33 - 0.27	0.30	0.02	
DOAE	0.29	0.32 - 0.25	0.29	0.02	0.30 - 0.25	0.28	0.01	
DFL	0.20	0.29 - 0.14	0.20	0.03	0.20 - 0.15	0.17	0.01	
DEAO	0.27	0.29 - 0.24	0.26	0.01	0.26 - 0.23	0.24	0.01	
AFL	0.12	0.12 - 0.09	0.11	0.01	0.11 - 0.07	0.09	0.01	
AEDE	0.17	0.19 - 0.16	0.17	0.01	0.20 - 0.16	0.17	0.01	
EDUP	0.23	0.28 - 0.23	0.26	0.02	0.29 - 0.21	0.24	0.03	
EDLP	0.28	0.33 - 0.28	0.30	0.01	0.33 - 0.26	0.30	0.02	
EAUP	0.29	0.32 - 0.28	0.30	0.01	0.31 - 0.26	0.29	0.01	
EALP	0.22	0.28 - 0.24	0.26	0.01	0.30 - 0.23	0.26	0.02	



**Figure 5.** (A), holotype, male (CPUM-1509) and (B), paratype, female (CPUM-1510) of *Zoogoneticus purhepechus* from La Luz Spring.

variation pattern within the populations of the 2 species analysed here show the separation of 2 well-defined groups, supporting previous findings made with molecular characters (Doadrio and Domínguez, 2004). The ANOVA analysis between populations shows that DOAE distance differs between them. We inferred that this is a result of the larger DFL distance in *Z. purhepechus* and this inference is supported by the PCA. In the same manner, differences in PrOL were obtained and supported by the minus preorbital length obtained as a diagnostic character in PCA. The morphometric diagnostic characters are preorbital length and dorsal fin length in the PCI. Accordingly, with this classification, which was corroborated by the DFA, the *Z. quitzeensis* populations (San Cristobal Spring and La Mintzita Spring in the Cuitzeo drainage and Zacapu Lake) have a larger superior mandible and a shorter base of the dorsal fin. On the other hand *Z. purhepechus* (La Luz Spring) exhibits a shorter upper mandible, a larger dorsal fin base and a higher number of dorsal rays.

This grouping model is congruent with the results of the Bayesian analysis, and the high genetic divergences between *Z. quitzeensis* and *Z. purhepechus* ( $D_{HKY} =$

**Table 6.** Molecular diagnostic characters for the cytochrome *b* gene in *Zoogoneticus* spp. (BPP = base pair position)

Species	BPP									
	136	237	327	481	585	798	816	870	883	895
<i>Z. tequila</i>	C	T	T	T	T	C	A	A	C	C
<i>Z. quitzeoensis</i>	T	T	T	G	A	A	A	C	C	C
<i>Z. purhepechus</i>	A	C	C	A	C	T	G	T	T	T

3.4%, range= 3-3.8%). These values are higher than those described for the family Goodeidae, in which an intraspecific pairwise uncorrected “p” distance of 0.001 to 1.7% was found, and are similar to those found in interspecific distances of 1.7 to 11% (Doadrio and Domínguez, 2004). In the same manner, the middle Lerma and lower Lerma basins are considered to have undergone a pattern of isolation and union, which correlates with vicariant events inferred to promote cladogenetic processes in at least 2 pairs of sister species within the Goodeinae (e.g. *Skiffia lermae*-*Skiffia multipunctata* and *Zoogoneticus quitzeoensis*-*Z. purhepechus*) (Domínguez-Domínguez et al., 2006a).

### Acknowledgments

The authors are grateful to Luis Escalera, Asdrubal Molina, Hugo Mejía, Rogelio Rosas, Jen Nightingale, Jean de Moree, Ludo Couvreur, Kees de Jong, Juan Carlos Merino and Adrian Pompa for their help during field trips. Partial funds to conduct this research were provided by the COECyT Michoacán and Chester Zoo Garden, England to OD and by the project CGL2006-12325/BOS. We thank Ivan Dibble for his support to the Hobbyist Aqualab Conservation Project (HALCP) that receives funds from European and North American aquarist associations. OD and RP thank the Consejo Nacional de Ciencia y Tecnología for the scholarship.

### Literature cited

- Barbour, C. D. 1973. A biogeographical history of *Chirostoma* (Pisces: Atherinidae): A species flock from the Mexican Plateau. *Copeia* 1973:533-556.
- Burnaby, T. P. 1966. Growth-invariant discriminant functions and generalized distances. *Biometrics* 22: 96-110.
- De la Vega-Salazar, M., E. Avila-Luna and C. Macías-García. 2003. Ecological evaluation of local extinction: the case of 2 genera of endemic Mexican fish, *Zoogoneticus* and *Skiffia*. *Biodiversity and Conservation* 12:2043-2056.
- Doadrio, I., J. A. Carmona and C. Fernández-Delgado. 2002. Morphometric study of the Iberian *Aphianus* (Actinopterygii, Cyprinodontiformes), with description of a new species. *Folia Zoologica* 51:67-79.
- Doadrio, I. and O. Domínguez. 2004. Phylogenetic relationships within the fish family Goodeidae based on cytochrome *b* sequence data. *Molecular Phylogenetics and Evolution* 31:416-430.
- SEMARNAT (Secretaría de Medio Ambiente y Recursos Naturales). 2002. Norma Oficial Mexicana NOM-059-ECOL-2001, Protección ambiental-Especies nativas de México de flora y fauna silvestres-Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-Lista de especies en riesgo. Diario Oficial de la Federación. 6 de marzo de 2002, Primera sección. México, DF.
- Domínguez-Domínguez, O., N. Mercado-Silva, J. Lyons and H. J. Grier. 2005. The viviparous Goodeid fish. *In* Proceedings of the second international symposium of viviparous fishes, M. C. Uribe and H. J. Grier (eds.). New Life Publications, Homestead, Florida. p. 505-549.
- Domínguez-Domínguez, O., I. Doadrio and G. Pérez-Ponce de León. 2006a. Historical biogeography of some river basins in Central Mexico evidenced by their goodeine freshwater fishes: A preliminary hypothesis using secondary Brooks Parsimony Analysis (BPA). *Journal of Biogeography* 30:1437-1447.
- Domínguez-Domínguez O., E. Martínez-Meyer, L. Zambrano and G. Pérez-Ponce de León. 2006b. Using ecological-niche modeling as a conservation tool for freshwater species: the live-bearing fishes (Goodeidae) in Central Mexico. *Conservation Biology*. 20: 1730-1739.
- Domínguez-Domínguez O., L. Boto, F. Alda, G. Pérez-Ponce de León and I. Doadrio. 2007. Human impacts on Basins of the Mesa Central of Mexico, and its genetic effects on an endangered fish, *Zoogoneticus quitzeoensis*. *Conservation Biology* 21:168-180.
- Echelle, A. A. and A. F. Echelle. 1984. Evolutionary genetics of a “species flock:” Atherinid fishes on the Mesa Central of Mexico. *In* Evolution of fish species flocks, A. A. Echelle and I. Kornfield (eds.). University of Maine Press, Orono. p. 93-110.
- Espinoza-Pérez, H., M. T. Gaspar-Dillanes and P. Fuentes-Mata. 1993. Listados faunísticos de México. III. Los peces dulceacuicolas mexicanos. Instituto de Biología, Universidad Nacional Autónoma de México, México, D.F. 98 pp.
- Grudzien, T. A., M. M. White and B. J. Turner. 1992. Biochemical systematics of the viviparous fish family Goodeidae. *Journal of Fish Biology* 40:810-814.
- Guzmán-Arroyo, F. 1994. Osteología y variación no geográfica

- de la suspensión de la aleta anal de *Goodea luitpoldi*, (Osteichthyes: Goodeidae). Universidad, Ciencia y Tecnología 3:33-41.
- Hubbs, C. L. and C. L. Turner. 1939. Studies of the fishes of the Order Cyprinodontes. XVI. A revision of the Goodeidae. Miscellaneous Publications of the Museum of Zoology, University of Michigan 42:1-80.
- Hueselsenbeck, J.P. and F. R. Ronquist. 2001. MrBayes: Bayesian inference of phylogeny. *Bioinformatics* 17:754-755.
- IUCN (International Union for the Conservation of Nature and Natural Resources). 1994. IUCN Red List Categories. Prepared by the IUCN Species Survival Commission. Gland: IUCN, Switzerland.
- Machordom, A. and I. Doadrio. 2001. Evidence of a Cenozoic Betic-Kabilian connection based on freshwater fish phylogeography (*Luciobarbus*, Cyprinidae). *Molecular Phylogenetics and Evolution* 18:252-263.
- Mateos, M., O. I. Sanjur and R. C. Vrijenhoek. 2002. Historical biogeography of the livebearing fish genus *Poeciliopsis* (Poeciliidae: Cyprinodontiformes). *Evolution* 56:972-984.
- Meek, S. E. 1904. The fresh-water fishes of Mexico north of the Isthmus of Tehuantepec. Field Columbian Museum Publications 5:1-252.
- Miller, R. R. and M. L. Smith. 1986. Origin and geography of the fishes of central Mexico. *In* The zoogeography of the North American Freshwater Fishes, C. H. Hocutt and E. O. Wiley (eds.). John Wiley and Sons, New York. p. 487-517.
- Parenti, L. 1981. A phylogenetic and biogeographic analysis of cyprinodontiform fishes (Teleostei, Atherinomorpha). *Bulletin of the American Museum of Natural History* 168:335-557.
- Posada, D. and K. A. Crandall. 1998. Modeltest: testing the model of DNA substitution. *Bioinformatics* 14: 817-818.
- Regan, C. T. 1908. Pisces. *In* *Biologia Centrali-Americana*, F. D. Godman and O. Salvin (eds.). London, United Kingdom. p. 1-203.
- Rohlf, F. J. and F. L. Bookstein. 1987. A comment on shearing as a method for size correction. *Systematic Zoology* 36:356-367.
- Rohlf, F. J. 2000. NTSYS-pc: Numerical taxonomy and multivariate analysis system, version 2.1. Setauket, New York.
- Sambrook, J., E. F. Fritsch and T. Maniatis. 1989. *Molecular Cloning: A Laboratory Manual*. New York: Cold Spring Harbor Laboratory.
- Schönhuth, M. S. and I. Doadrio. 2003. Phylogenetic relationships of Mexican minnows of the genus *Notropis* (Actinopterygii, Cyprinidae). *Biological Journal of the Linnean Society* 80:323-337.
- Webb, S. A. and R. R. Miller. 1998. *Zoogoneticus tequila* a new goodeid fish (Cyprinodontiformes) from Ameca drainage of Mexico, and a rediagnosis of the genus. *Occasional Papers of the Museum of Zoology, University of Michigan* 725:1-23.